

A Comparative Analysis of Lateral Suspension Techniques and Sacrocolpopexy for Pelvic Organ Prolapse Repair in a Tertiary Care Center

Pelvik Organ Prolapsusu Cerrahisinde Üçüncü Basamak Bir Merkezde Yapılan Sakrokolpopeksi ile Lateral Süspansiyon Tekniklerinin Karşılaştırılması

Deniz Taşkiran¹, Sadettin Oğuzhan Tutar¹, Salih Kolsuz¹, Öznur Tosun²

¹Giresun University Faculty of Medicine, Department of Obstetrics and Gynecology, Giresun, Türkiye

²University of Health Sciences Türkiye, Antalya City Hospital, Clinic of Gynecology and Obstetrics, Antalya, Türkiye

ABSTRACT

Background: Pelvic organ prolapse (POP) is a common condition in elderly women and has an enormous quality-of-life impact. Various surgical techniques, including vaginal, abdominal, open (laparotomy), and laparoscopic approaches, are utilized based on patient symptoms and clinical presentation. This study aimed to compare the perioperative outcomes and clinical results of laparotomic sacrocolpopexy (L-SCP), considered the gold standard for POP repair, with those of laparoscopic lateral suspension (LLS).

Materials and Methods: A retrospective analysis was conducted of 186 POP surgeries performed at a tertiary care center between 2022 and 2024. Patients who met the inclusion criteria were divided into two groups. 24 of these patients underwent L-SCP, and 25 underwent LLS. The parameters compared included age, body mass index, operative time, length of postoperative hospital stay, perioperative hemoglobin change, postoperative POP stage, and recurrence and complication rates. The primary outcome was the objective anatomical success rate, assessed using the POP Quantification system.

Results: The mean operative time was significantly longer in the L-SCP group than in the LLS group. Duration of hospital stay and perioperative hemoglobin changes were similar in the two groups. Although the earlier stages of preoperative prolapse were more common in the L-SCP group, anatomical outcomes were comparable at 12 months postoperatively. In the L-SCP group, the complications included one case (4.1%) of common iliac vein injury, one case (4.1%) of ureteral injury, and three cases (12.5%) of postoperative surgical site infection. In the LLS group, there were two cases (8%) of bladder injury, one case (4%) of ureteral injury, two cases (8%) of mesh-related inflammatory reaction, and one case (4%) of surgical site infection.

Conclusion: Laparotomic and laparoscopic approaches showed similar short-term anatomical outcomes and recurrence rates. Laparotomic SCP was associated with longer operative time and higher infection risk, while laparoscopic LS had a different complication profile. The surgical approach should be guided by patient characteristics and the surgeon's experience, rather than by presumed superiority.

Keywords: Pelvic organ prolapse, sacrocolpopexy, lateral suspension

ÖZ

Amaç: Pelvik organ prolapsusu (POP), özellikle ileri yaş kadınlarda sık görülen ve yaşam kalitesini düşüren klinik bir durumdur. Hastaların semptom ve şikayetlerine göre farklı tekniklerle hastalara uygun cerrahi yapılmaktadır. Vajinal, abdominal, laparotomik ya da laparoskopik cerrahi teknikler kullanılabilir. POP'da altın standart teknik olan laparotomik sakrokolpopeksi (L-SCP) ile son yıllarda popüler olan laparoskopik lateral süspansiyon (LLS) yapılan hastalar karşılaştırılmıştır.

Gereç ve Yöntemler: 2022–2024 yılları arasında üçüncü basamak bir hastanede gerçekleştirilen 186 POP cerrahi vakaları retrospektif olarak değerlendirildi. Yirmidört tane L-SCP ve 25 tane LLS cerrahi teknik olmak üzere iki gruba ayrıldı. Yaş, vücut kitle indeksi, operasyon süresi, postoperatif hastanede kalış süresi, perioperatif hemoglobinin değişimi, postoperatif prolapsus evresi, nüks sayısı ve komplikasyon oranları karşılaştırıldı. Çalışmanın birincil sonucu olarak Pelvik Organ Prolapsusu Kantifikasyon sistemine göre objektif kür sağlama oranı değerlendirildi.



Address for Correspondence: Deniz Taşkiran, Giresun University Faculty of Medicine, Department of Obstetrics and Gynecology, Giresun, Türkiye

E-mail: deniztaskiran93@gmail.com **ORCID ID:** orcid.org/0000-0001-7270-4566

Received: 30.10.2025 **Accepted:** 25.02.2026 **Epub:** 22.05.2026

Cite this article as: Taşkiran D, Tutar SO, Kolsuz S, Tosun Ö. A comparative analysis of lateral suspension techniques and sacrocolpopexy for pelvic organ prolapse repair in a tertiary care center. Hamidiye Med J.



Bulgular: Ameliyat süresi L-SCP grubunda, LLS grubuna göre daha yüksekti. Her iki grupta hospitalizasyon süresi benzerdi. Hastaların perioperatif hemoglobin değişimleri her iki grupta benzerdi. Preoperatif desensus seviyesi L-SCP grubunda daha yüksekti. Ancak ameliyat sonrası 12 aylık desensus seviyesi her iki grupta benzerdi. L-SCP grubunda 1 hastada (%4,1) common iliak ven yaralanması, 1 hastada (%4,1) üreter yaralanması, 3 hastada (%12,5) post operatif cerrahi alan enfeksiyonu görüldü. LLS grubunda 2 hastada (%8) mesane yaralanması, 1 hastada (%4) üreter hasarı, 2 hastada (%8) meshe bağlı inflamatuvar reaksiyonu ve 1 hastada (%4) cerrahi alan enfeksiyonu görüldü.

Sonuç: L-SCP ve laparoskopik lateral süspansiyon kısa dönem anatomik sonuçlar ve nüks oranları açısından benzer gözlemsel sonuçlar göstermiştir. Bununla birlikte, cerrahi yaklaşım farklılığı ve gruplar arasındaki preoperatif prolapsus şiddeti eşitsizliği nedeniyle bu bulgular dikkatle yorumlanmalıdır. L-SCP daha uzun ameliyat süresi ve enfeksiyon riski ile ilişkililikten, LLS farklı bir komplikasyon profili sergilemiştir. Bu nedenle cerrahi teknik seçimi, yöntemlerin mutlak üstünlüğü yerine hasta özellikleri ve cerrah deneyimi doğrultusunda bireyselleştirilmelidir.

Anahtar Kelimeler: Pelvik organ prolapsusu, sakrokolpopeksi, lateral süspansiyon

Introduction

Pelvic organ prolapse (POP) is a clinical syndrome in which pelvic organs sag into or below the vaginal canal, typically resulting in vaginal bulging and pelvic pressure. It results primarily from a defect or weakness within the connective tissue and supporting structures of the pelvic floor. POP most often occurs in women aged 45–85 years (1). It has been estimated that nearly 13% of women will require surgical correction for POP at some stage in their lives (2). As life expectancy increases, the prevalence of symptomatic POP among women is projected to rise to 46% by 2050 (3).

Although several factors associated with POP have been ascertained, the precise associations between these factors and established risk factors remain to be identified (4). Known risk factors include menopause, increased parity, obesity, advanced age, genetic predisposition, and connective tissue disorders.

Diagnosis of prolapse is based on patient history and physical examination, with the physical examination assessing the anterior vaginal wall, vaginal apex, and posterior vaginal wall. POP staging is performed based on these findings, typically using the POP Quantification (POP-Q) system. When surgery is indicated, the surgical approach (transvaginal, laparoscopic, or open laparotomy) and the choice between native-tissue repair and mesh use are determined by the patient's overall health status and specific characteristics. The vaginal route is less invasive and is associated with lower morbidity than laparotomy and laparoscopy (5).

In recent years, increasing complication rates associated with mesh use have been reported (6). These complications include vaginal mucosal erosion, pain, infection, prolapse recurrence, and bladder or bowel injury during mesh implantation (7). Even in carefully selected patients,

judicious use of mesh prevents recurrence of prolapse and improves long-term quality of life by ensuring pelvic floor support (8). Effective treatment of POP development and postoperative recurrence requires careful preoperative counseling, understanding patient expectations, and individualized surgical planning (9).


Abdominal sacrocolpopexy (SCP) involves attaching the vaginal apex to the anterior longitudinal ligament using mesh (10). This procedure, described by DeLancey (11), is indicated for all POP patients with level 1 defects. In young, sexually active women, SCP remains the gold standard for apical prolapse repair. However, data on its short- and long-term success rates and complication rates are conflicting. SCP demands extensive pelvic dissection and advanced suturing skills. To mitigate the risk of complications, a less complex lateral suspension (LS) technique was developed. LS is performed by placing a T-shaped mesh in the vesicovaginal space and securing it with sutures to the anterior vaginal wall, cervix, and isthmus (12). Over the past decade, multiple studies have assessed the efficacy and complication profile of this technique.

The present study aims to compare the short-term outcomes of abdominal SCP, the gold standard for POP and apical prolapse repair, with those of laparoscopic LS, a technically less demanding procedure with a potentially lower risk of complications.

Materials and Methods

A retrospective review was conducted of 186 POP cases performed at a tertiary care hospital between January 2022 and January 2024. Twelve cases were excluded due to vaginal cuff prolapse, 42 due to isolated anterior colporrhaphy, 24 due to isolated posterior colporrhaphy, and 59 due to incomplete data (Figure 1).

Due to the limited number of surgeons experienced in performing laparoscopic SCP at our center, various



suspension techniques were compared across different surgical approaches.

The final analysis included 25 patients who underwent laparoscopic LS with hysterectomy and 24 who underwent open SCP with hysterectomy, constituting a retrospective cohort. Patients who underwent a hysterectomy planned for prolapse were included in the study. Patients who underwent a hysterectomy for myoma uteri or other indications were excluded.

The demographic and clinical variables compared included age, operative time, body mass index (BMI), postoperative hospital stay, postoperative change in hemoglobin, postoperative prolapse stage, and recurrence and complication rates. Patients with preoperative apical prolapse stage 2 or higher, according to the POP-Q system, were included. Recurrences were defined as stage III or higher prolapse occurring within one year postoperatively. The primary outcome was the objective cure rate evaluated by the POP-Q system.

The study protocol was approved by the Scientific Research Ethics Committee of Giresun Training and Research Hospital (decision number: 30.10.2024/04, dated: 04.11.2024). All procedures were conducted in accordance with the Ethical Principles for Medical and Health Research Involving Human Subjects.

Statistical Analysis

All statistical analyses were performed using SPSS version 22.0 (IBM Corp., Armonk, NY, USA). The normality of continuous variables was assessed using the Shapiro–Wilk test. Continuous variables that were not normally distributed

were summarized as medians and interquartile ranges while categorical variables were presented as numbers and percentages (n, %). Comparisons between the two groups were performed using the Mann–Whitney U test for non-normally distributed continuous variables and the chi-square test or Fisher’s exact test for categorical variables, as appropriate. Preoperative and postoperative POP-Q stages were treated as ordinal variables and summarized using the median (range) and frequencies for each stage. A p-value <0.05 was considered statistically significant. Due to the retrospective nature of the study and the small sample size, no formal post hoc power analysis was performed; however, the effect sizes were calculated where relevant to assess clinical relevance.

Results

This retrospective cohort study included 49 patients with POP who underwent either laparotomic SCP (L-SCP) (n = 24) or laparoscopic LS (n = 25), all of whom underwent concomitant hysterectomy (Figure 1). Baseline characteristics, including age (median 60.5 vs. 63.5 years), duration of menopause, BMI, gravidity, parity, and comorbidities such as hypertension and diabetes, did not differ significantly (all p > 0.05). Operative time was significantly shorter in the LS group than in the SCP group (median 150 vs. 177.5 minutes, p = 0.002), reflecting the minimally invasive nature of the laparoscopic approach. Preoperative hemoglobin levels were slightly higher in SCP patients (13.45 vs. 12.65 g/dL, p = 0.033), and postoperative values were also higher (11.95 vs. 11.20 g/dL, p = 0.012); however, the magnitude of the hemoglobin drop and transfusion rates did not differ significantly (Table 1).

Preoperative POP-Q stage distribution differed significantly between groups, with more advanced prolapse (Stages 3–4) observed in the SCP group ($p = 0.027$). At 12 months postoperatively, the distribution of POP-Q stages was comparable between the two groups, with no statistically significant difference. In the postoperative assessment, recurrences of the apical prolapse were evaluated together with those of the anterior and posterior compartments (Table 2).

Complication profiles were comparable, though bladder injury (8%) and mesh-related inflammatory reaction (8%)

occurred exclusively in the LS group, whereas vascular injury (4.1%) and higher surgical-site infection rates (12.5%) occurred in the SCP group. Length of hospital stay and rates of injury to adjacent organs were similar between the groups (Table 3). Collectively, these data support the conclusion that both SCP and LS are effective and safe for POP correction, with the laparoscopic approach associated with reduced operative time and slightly lower perioperative blood loss without increasing the risk of complications.

Table 1. Comparison between groups.

	L-SCP (n = 24)	Laparoscopic LS (n = 25)	p-value
Age (years), median (IQR)	60.5 (54–67)	63.5 (55–71)	0.441
Menopause Duration (years), median (IQR)	12 (6–18)	14 (7–21)	0.507
BMI (kg/m ²), median (IQR)	26.15 (24–31)	28.3 (25–31)	0.395
Gravidity, median (IQR), median (IQR)	3 (2–4)	4 (2–5)	0.975
Parity, median (IQR)	2 (2–3)	3 (2–4)	0.819
Normal vaginal delivery, median (IQR)	2.5 (2–3)	3 (2–4)	0.758
Cesarean delivery, median (IQR)	0 (0–0)	0 (0–0)	0.584
Operative time (minutes), median (IQR)	177.5 (160–210)	150.00 (120–180)	0.002*
Hospital stay (days), median (IQR)	3 (3–4)	3 (3–4)	0.078
Preoperative hemoglobin (g/dL), median (IQR)	13.45 (12.5–14.2)	12.65 (11.9–13.4)	0.033*
Postoperative hemoglobin (g/dL), median (IQR)	11.95 (10.8–12.6)	11.20 (10.5–12.0)	0.012*
Hemoglobin change (g/dL), median (IQR)	1.45 (1.0–2.0)	1.3 (1.0–2.1)	0.489
Blood transfusion, n (%)	1 (4.1%)	3 (12%)	0.301
Adjacent organ injury, n (%)	2 (8.3%)	3 (12%)	0.673
Hypertension, n %	14 (58.3%)	11 (44%)	0.199
Diabetes mellitus, n (%)	10 (41.6%)	12 (48%)	0.880
Smoking, n (%)	9 (37.5%)	6 (24%)	0.474

Data are presented as median (IQR) or number (%). p-values calculated using Mann–Whitney U test for continuous variables and Fisher’s exact test for categorical variables. Statistically significant p-values are shown in bold. BMI, body mass index; IQR, interquartile range; L-SCP, laparotomic sacrocolpopexy; LS, lateral suspension; n, number of patients.

Table 2. Preoperative and 12-month postoperative POP-Q stages in L-SCP and laparoscopic LS groups.

POP-Q stage	Laparotomic SCP (n = 24)	Laparoscopic LS (n = 25)	p-value
Preoperative POP-Q stage			
Stage 1	0 (0%)	0 (0%)	0.027*
Stage 2	1 (4.1%)	1 (4.0%)	
Stage 3	10 (41.6%)	17 (68.0%)	
Stage 4	13 (54.1%)	7 (28.0%)	
Postoperative POP-Q stage (12 months)			
Stage 1	14 (58.3%)	12 (48.0%)	0.196
Stage 2	8 (33.3%)	9 (36.0%)	
Stage 3	1 (4.1%)	2 (8.0%)	
Stage 4	1 (4.1%)	1 (4.0%)	

p-values calculated using Fisher’s exact test. L-SCP, laparotomic sacrocolpopexy; LS, lateral suspension; n, number of patients; POP-Q, Pelvic Organ Prolapse Quantification.

Table 3. Postoperative complications and perioperative outcomes.

	Laparotomic SCP (n = 24)	Laparoscopic LS (n = 25)
Vascular injury (common iliac vein)	1(4.1%)	0 (0%)
Bladder injury	0 (0%)	2 (8%)
Ureteral damage	1 (4.1%)	1 (4%)
Mesh-related inflammatory reaction	0 (0%)	2 (8%)
Surgical site infection	3 (12.5%)	1 (4%)

LS, lateral suspension; n, number of patients; SCP, sacrocolpopexy.

Discussion

The clinical management of POP remains challenging, with varying success rates reported among different surgical approaches. Minimally invasive techniques have gained popularity because they reduce operative morbidity, shorten hospital stays, and accelerate recovery. In our study, LS demonstrated significantly shorter operative times than abdominal SCP, consistent with the literature (13,14). Despite these differences, 12-month postoperative prolapse descent was comparable between LS and SCP, regardless of preoperative POP severity. Several studies support the efficacy of LS. One series reported 92% anterior compartment and 100% apical compartment recovery at 12 months post-laparoscopic LS (13). For an inexperienced surgeon, laparoscopic LS takes approximately 184 minutes (14). Learning-curve analyses indicate that LS is technically less demanding than laparoscopic SCP, while achieving similar objective success at 12 months (LS 89.1% vs. SCP 90.7%) (14). Laparoscopic LS is a less complex procedure to learn, and its success rate matches that of SCP. The degree of descent was the same for both procedures when compared at the 12-month postoperative follow-up in our study. Laparoscopic LS and SCP have equal efficacy and success rates in the management of POP (15). In another study, the primary cure rate of apical prolapse after laparoscopic LS was 87.5%, and after laparoscopic SCP was 100%. Laparoscopic LS, as a procedure with a low-risk of complications and short-term objective outcomes, may be an alternative to laparoscopic SCP (16). In our study, both techniques demonstrated similar effectiveness and short-term success. Our finding that LS was associated with shorter operative time and higher 12-month success rate is fully consistent with the literature. While some literature favors SCP for apical support, your data demonstrate that LS does not compromise anatomical success despite its technical simplicity. While SCP remains a robust technique for advanced prolapse, LS offers a shorter operative duration and comparable short-term objective success. Given its shorter learning curve and equivalent efficacy at 1 year, LS

should be considered a first-line minimally invasive option, particularly in surgical settings where reducing anesthesia time and technical complexity is prioritized. LS is a strong, reliable, and minimally invasive alternative to SCP, with its advantages in learning curve and operative time.

Comparing abdominal SCP and abdominal LS, apical prolapse repair rates were 94% and 92%, respectively, with both being equally effective at 12-month follow-up for severe apical prolapse (17). Anatomical success rates of 81.8% for SCP and 90% for LS were reported in a series of 93 patients; the treatments were considered equally effective (18). In our cohort, two patients in the SCP group and three patients in the LS group developed stage 3–4 prolapse at 12 months postoperatively, underscoring comparable success between the groups in managing advanced apical prolapse. The success rates in the literature parallel our low recurrence counts: 2 patients in SCP and 3 patients in LS. The primary finding is that the minimally invasive nature of LS does not compromise anatomical success. Since both techniques show similar efficacy in advanced apical prolapse, the choice should be based on logistical considerations and safety, rather than on success rates. The choice between SCP and LS may be guided more by surgical logistics and perioperative safety profiles than by differences in anatomical success, as both procedures offer high efficacy for long-term management of prolapse.

Pelvic dissection, particularly in the presacral region, carries inherent risks due to the proximity of critical structures. Previous studies report a 1.0% risk of ureteral injury and a 4.4% risk of vascular damage during pelvic surgery, with complications potentially exacerbated by anatomical variations, such as the more medial course of the left iliac vessels (19). Consistent with this, in our SCP cohort, vascular and ureteral injuries occurred in 4.1% of patients, highlighting the inherent intraoperative risks of the open approach. Wound-related complications, including surgical site infections and intra-abdominal abscesses, are also well-documented, with reported rates ranging from 2–12.5% for infections and 8% for abscesses (20). Bladder injury appears less common; a meta-analysis found no

significant difference in bladder perforation rates between obese and non-obese patients undergoing SCP (21), and similarly, no bladder perforation was observed in non-obese women in our series. Surgical wound infection rates are typically lower with laparoscopy because tissue damage is reduced and immune function is preserved (22). Our findings corroborate this, as the laparoscopic LS group demonstrated lower surgical site infection rates than SCP, supporting the advantages of minimally invasive techniques. Beyond surgical technique, patient-related factors also influence POP outcomes. Hormonal changes in menopause reduce estrogen levels, which, in turn, alter the collagen content and compromise pelvic tissue strength (23). Additionally, comorbidities such as diabetes mellitus and hypertension contribute to POP pathogenesis (24). In our study, a similar prevalence of these comorbidities in both groups suggests that the observed differences in complications were largely attributable to the surgical approach rather than to patient baseline characteristics. Overall, these findings emphasize the importance of surgical technique in minimizing perioperative complications and highlight the role of patient factors in POP management. Laparoscopic LS appears to offer a safer perioperative profile without compromising anatomical outcomes, particularly in patients with comparable comorbidity profiles.

Laparoscopic LS provides a safer perioperative profile than L-SCP, with lower rates of surgical site infections and major vascular or ureteral injuries. Both techniques achieve comparable 12-month anatomical success, making LS a highly effective and less invasive alternative for POP management. While LS is technically less demanding, it carries a distinct complication profile, including a higher risk of bladder injury and mesh-related inflammatory reactions, whereas SCP carries inherent risks of vascular and ureteral injury. The lower infection rates with LS and a similar distribution of comorbidities suggest that these outcomes primarily reflect the surgical approach rather than patient-related factors.

Study Limitations

This study has several important limitations. Its retrospective design and relatively small sample size restrict causal inference and the generalizability of the findings. The comparison of an open surgical approach (L-SCP) with a minimally invasive technique (laparoscopic LS) introduces potential confounding related to surgical access, rather than to the suspension method itself. Additionally, the limited number of experienced surgeons performing laparoscopic SCP at our center necessitated using different surgical approaches for the various suspension techniques, which represents a notable limitation. Preoperative hemoglobin

levels differed significantly between groups; however, perioperative hemoglobin changes were analyzed without adjustment. This factor should be considered when interpreting the results and acknowledged as a limitation of the study.

Conclusion

In this retrospective cohort study, L-SCP and laparoscopic LS demonstrated comparable short-term anatomical outcomes and recurrence rates for POP surgery. L-SCP was associated with longer operative time and higher infectious morbidity, whereas laparoscopic LS showed a different complication profile, including mesh-related events. Given the methodological limitations and differences in surgical approach, no definitive superiority can be inferred. Surgical technique selection should, therefore, be individualized based on patient characteristics, disease severity, and the surgeon's experience. Prospective, approach-matched studies are needed to further clarify comparative effectiveness.

Ethics

Ethics Committee Approval: The study protocol was approved by the Scientific Research Ethics Committee of Giresun Training and Research Hospital (decision number: 30.10.2024/04, dated: 04.11.2024).

Informed Consent: Retrospective cohort study.

Footnotes

Authorship Contributions

Surgical and Medical Practices: D.T., S.K., Ö.T., Concept: D.T., S.O.T., Design: D.T., S.K., Data Collection or Processing: D.T., S.O.T., Ö.T., Analysis or Interpretation: D.T., S.K., Literature Search: D.T., S.O.T., Ö.T., Writing: D.T., S.O.T., Ö.T.

Conflict of Interest: No conflict of interest was declared by the author(s).

Financial Disclosure: The author(s) declared that this study received no financial support.

REFERENCES

1. Weintraub AY, Gliner H, Marcus-Braun N. Narrative review of the epidemiology, diagnosis and pathophysiology of pelvic organ prolapse. *Int Braz J Urol.* 2019;46:5-14. [[Crossref](#)]
2. Raju R, Linder BJ. Evaluation and management of pelvic organ prolapse. *Mayo Clin Proc.* 2021;96:3122-3129. [[Crossref](#)]
3. Wu JM, Hundley AF, Fulton RG, Myers ER. Forecasting the prevalence of pelvic floor disorders in U.S. Women: 2010 to 2050. *Obstet Gynecol.* 2009;114:1278-1283. [[Crossref](#)]
4. Petros PE, Ulmsten UI. An integral theory of female urinary incontinence. Experimental and clinical considerations. *Acta Obstet Gynecol Scand Suppl.* 1990;153:7-31. [[Crossref](#)]

5. Maher C, Yeung E, Haya N, Christmann-Schmid C, Mowat A, Chen Z, et al. Surgery for women with apical vaginal prolapse. *Cochrane Database Syst Rev.* 2023;7:CD012376. [[Crossref](#)]
6. Dibb B, Woodgate F, Taylor L. When things go wrong: experiences of vaginal mesh complications. *Int Urogynecol J.* 2023;34:1575-1581. [[Crossref](#)]
7. Kasyan G, Abramyan K, Popov AA, Gvozdev M, Pushkar D. Mesh-related and intraoperative complications of pelvic organ prolapse repair. *Cent European J Urol.* 2014;67:296-301. [[Crossref](#)]
8. Maher C, Feiner B, Baessler K, Schmid C. Surgical management of pelvic organ prolapse in women. *Cochrane Database Syst Rev.* 2013:CD004014. Update in: *Cochrane Database Syst Rev.* 2016;11:CD004014. [[Crossref](#)]
9. Salvatore S, Siesto G, Serati M. Risk factors for recurrence of genital prolapse. *Curr Opin Obstet Gynecol.* 2010;22:420-424. [[Crossref](#)]
10. Visco AG, Advincula AP. Robotic gynecologic surgery. *Obstet Gynecol.* 2008;112:1369-1384. [[Crossref](#)]
11. DeLancey JO. Anatomic aspects of vaginal eversion after hysterectomy. *Am J Obstet Gynecol.* 1992;166:1717-1728. [[Crossref](#)]
12. Campagna G, Vacca L, Panico G, Caramazza D, Lombisani A, Scambia G, et al. Laparoscopic lateral suspension for pelvic organ prolapse: a systematic literature review. *Eur J Obstet Gynecol Reprod Biol.* 2021;264:318-329. [[Crossref](#)]
13. Martinello R, Scutiero G, Stuto A, Indraccolo U, Cracco F, Borghi C, et al. Correction of pelvic organ prolapse by laparoscopic lateral suspension with mesh: a clinical series. *Eur J Obstet Gynecol Reprod Biol.* 2019;240:351-356. [[Crossref](#)]
14. Malanowska-Jarema E, Osnytska Y, Starczewski A, Balzarro M, Rubilotta E. A comparative study in learning curves of laparoscopic lateral suspension vs. laparoscopic sacrocolpopexy: preliminary results. *Front Surg.* 2023;10:1274178. [[Crossref](#)]
15. Bueno Garcia Reyes P, Hashim H. Mesh complications: best practice in diagnosis and treatment. *Ther Adv Urol.* 2020;12:1756287220942993. [[Crossref](#)]
16. Isenlik BS, Aksoy O, Erol O, Mulayim B. Comparison of laparoscopic lateral suspension and laparoscopic sacrocolpopexy with concurrent total laparoscopic hysterectomy for the treatment of pelvic organ prolapse: a randomized controlled clinical trial. *Int Urogynecol J.* 2023;34:231-238. [[Crossref](#)]
17. Russo E, Montt Guevara MM, Sacinti KG, Misasi G, Falcone M, Morganti R, et al. Minimal invasive abdominal sacral colpopexy and abdominal lateral suspension: a prospective, open-label, multicenter, non-inferiority trial. *J Clin Med.* 2023;12:2926. [[Crossref](#)]
18. Malanowska-Jarema E, Starczewski A, Melnyk M, Oliveira D, Balzarro M, et al. A randomized clinical trial comparing dubuisson laparoscopic lateral suspension with laparoscopic sacropexy for pelvic organ prolapse: short-term results. *J Clin Med.* 2024;13:1348. [[Crossref](#)]
19. Nygaard IE, McCreery R, Brubaker L, Connolly A, Cundiff G, Weber AM, et al. Abdominal sacrocolpopexy: a comprehensive review. *Obstet Gynecol.* 2004;104:805-823. [[Crossref](#)]
20. Sato H, Abe H, Ikeda A, Miyagawa T, Sato K. Complications and clinical outcomes of laparoscopic sacrocolpopexy for pelvic organ prolapse. *J Obstet Gynaecol.* 2021;41:128-132. [[Crossref](#)]
21. Wen Q, Zhao Z, Wen J, Yang Y, Wang L, Wu J, et al. Impact of obesity on operative complications and outcome after sacrocolpopexy: a systematic review and meta-analysis. *Eur J Obstet Gynecol Reprod Biol.* 2021;258:309-316. [[Crossref](#)]
22. Targarona EM, Balagué C, Knook MM, Triás M. Laparoscopic surgery and surgical infection. *Br J Surg.* 2000;87:536-544. [[Crossref](#)]
23. Jackson SR, Avery NC, Tarlton JF, Eckford SD, Abrams P, Bailey AJ. Changes in metabolism of collagen in genitourinary prolapse. *Lancet.* 1996;347:1658-1661. [[Crossref](#)]
24. Isik H, Aynioglu O, Sahbaz A, Selimoglu R, Timur H, Harma M. Are hypertension and diabetes mellitus risk factors for pelvic organ prolapse? *Eur J Obstet Gynecol Reprod Biol.* 2016;197:59-62. [[Crossref](#)]